

## Lower 6 Chapter 10

# Trigonometric Identities and Equations

## Chapter Overview

1. Know exact trig values for  $30^\circ$ ,  $45^\circ$ ,  $60^\circ$  and understand unit circle.
2. Use identities  $\frac{\sin x}{\cos x} \equiv \tan x$  and  $\sin^2 x + \cos^2 x \equiv 1$
3. Solve equations of the form  $\sin(n\theta) = k$  and  $\sin(\theta \pm \alpha) = k$
4. Solve equations which are quadratic in  $\sin/\cos/\tan$ .

**Trigonometry**

5.3	Understand and use $\tan \theta = \frac{\sin \theta}{\cos \theta}$  Understand and use $\sin^2 \theta + \cos^2 \theta = 1$	These identities may be used to solve trigonometric equations or to prove further identities.
5.4	Solve simple trigonometric equations in a given interval, including quadratic equations in sin, cos and tan and equations involving multiples of the unknown angle.	Students should be able to solve equations such as $\sin(x + 70^\circ) = 0.5$ for $0 < x < 360^\circ$ , $3 + 5 \cos 2x = 1$ for $-180^\circ < x < 180^\circ$ $6 \cos^2 x^\circ + \sin x^\circ - 5 = 0$ , $0 \leq x < 360^\circ$ giving their answers in degrees.


## sin/cos/tan of 30°, 45°, 60°

You will frequently encounter angles of 30°, 60°, 45° in geometric problems. Why?


Although you will always have a calculator, you need to know how to derive these.

**All you need to remember:**

** Draw half a unit square and half an equilateral triangle of side 2.**




$$\begin{aligned}\sin(45^\circ) &= \\ \cos(45^\circ) &= \\ \tan(45^\circ) &= \end{aligned}$$



$$\begin{aligned}\sin(30^\circ) &= \\ \cos(30^\circ) &= \\ \tan(30^\circ) &= \\ \sin(60^\circ) &= \\ \cos(60^\circ) &= \\ \tan(60^\circ) &= \end{aligned}$$

## The Unit Circle and Trigonometry

For values of  $\theta$  in the range  $0 < \theta < 90^\circ$ , you know that  $\sin \theta$  and  $\cos \theta$  are lengths on a right-angled triangle:





The unit circle explains the behaviour of the trigonometric graphs beyond  $90^\circ$ .

However, the easiest way to remember whether  $\sin(x)$ ,  $\cos(x)$ ,  $\tan(x)$  are positive or negative is to just do a **very quick sketch (preferably mentally!)** of the corresponding graph.

**Note:** The textbook uses something called '*CAST diagrams*'. We will not be using them in this booklet, but you may wish to look at this technique as an alternative approach to various problems in the chapter.


### A Few Trigonometric Angle Laws

The following are all easily derivable using a quick sketch of a trigonometric graph, and are merely a convenience so you don't always have to draw out a graph every time.

You are highly encouraged to **memorise these** so that you can do exam questions faster.

1  $\sin(x) = \sin(180^\circ - x)$


e.g.  $\sin(150^\circ) = \sin(30^\circ)$



We saw this in the previous chapter when covering the 'ambiguous case' when using the sine rule.


2  $\cos(x) = \cos(360^\circ - x)$

e.g.  $\cos(330^\circ) = \cos(30^\circ)$



3 *sin* and *cos* repeat every  $360^\circ$  but *tan* every  $180^\circ$

e.g.  $\sin(390^\circ) = \sin(30^\circ)$



4  $\sin(x) = \cos(90^\circ - x)$

e.g.  $\sin(50^\circ) = \cos(40^\circ)$

Remember from the previous chapter that "cosine" by definition is the sine of the "complementary" angle. This was/is never covered in the textbook but caught everyone by surprise when it came up in a C3 exam.

## Examples

Without a calculator, work out the value of each below:

$$\tan(225^\circ) =$$

$$\tan(210^\circ) =$$

$$\sin(150^\circ) =$$

$$\cos(300^\circ) =$$

$$\sin(-45^\circ) =$$

$$\cos(750^\circ) =$$

$$\cos(120^\circ) =$$

$$\cos(315^\circ) =$$

$$\sin(420^\circ) =$$

$$\tan(-120^\circ) =$$

$$\tan(-45^\circ) =$$

$$\sin(135^\circ) =$$

## Trigonometric identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\sin^2 \theta + \cos^2 \theta = 1$$

### Examples

Prove that  $1 - \tan \theta \sin \theta \cos \theta \equiv \cos^2 \theta$

Prove that  $\tan \theta + \frac{1}{\tan \theta} \equiv \frac{1}{\sin \theta \cos \theta}$

Simplify  $5 - 5 \sin^2 \theta$



Test your understanding

Prove that  $\frac{\tan x \cos x}{\sqrt{1-\cos^2 x}} \equiv 1$

Prove that  $\frac{\cos^4 \theta - \sin^4 \theta}{\cos^2 \theta} \equiv 1 - \tan^2 \theta$

Prove that  $\tan^2 \theta \equiv \frac{1}{\cos^2 \theta} - 1$

Exercise 10C Pg 211/212

**Extension:**

[MAT 2008 1C] The simultaneous equations in  $x, y$ ,

$$(\cos \theta)x - (\sin \theta)y = 2$$

$$(\cos \theta)x + (\sin \theta)y = 1$$

are solvable:

- A) for all values of  $\theta$  in range  $0 \leq \theta < 2\pi$
- B) except for one value of  $\theta$  in range  $0 \leq \theta < 2\pi$
- C) except for two values of  $\theta$  in range  $0 \leq \theta < 2\pi$
- D) except for three values of  $\theta$  in range  $0 \leq \theta < 2\pi$

### Solving trigonometric equations

Solve  $\sin \theta = \frac{1}{2}$  in the interval  $0 \leq \theta \leq 360^\circ$ .

Solve  $5 \tan \theta = 10$  in the interval  $-180^\circ \leq \theta < 180^\circ$

Solve  $\sin \theta = -\frac{1}{2}$  in the interval  $0 \leq \theta \leq 360^\circ$ .

Solve  $\sin \theta = \sqrt{3} \cos \theta$  in the interval  $0 \leq \theta \leq 360^\circ$ .

Solve  $2 \cos \theta = \sqrt{3}$  in the interval  $0 \leq \theta \leq 360^\circ$ .

Solve  $\sqrt{3} \sin \theta = \cos \theta$  in the interval  $-180^\circ \leq \theta \leq 180^\circ$ .

### Harder equations

Solve  $\cos 3x = -\frac{1}{2}$  in the interval  $0 \leq x \leq 360^\circ$ .

Solve  $\sin(2x + 30^\circ) = \frac{1}{\sqrt{2}}$  in the interval  $0 \leq x \leq 360^\circ$ .

Solve  $\sin x = 2 \cos x$  in the interval  $0 \leq x < 300^\circ$

Solve, for  $0 \leq x < 180^\circ$ ,

$$\cos(3x - 10^\circ) = -0.4,$$

giving your answers to 1 decimal place. You should show each step in your working.

(7)

### Quadratics in sin/cos/tan

Solve  $5 \sin^2 x + 3 \sin x - 2 = 0$  in the interval  $0 \leq x \leq 360^\circ$ .

Solve  $\tan^2 \theta = 4$  in the interval  $0 \leq x \leq 360^\circ$ .



Solve  $2 \cos^2 x + 9 \sin x = 3 \sin^2 x$  in the interval  $-180^\circ \leq x \leq 180^\circ$ .

(a) Show that the equation

$$5 \sin x = 1 + 2 \cos^2 x$$

can be written in the form

$$2 \sin^2 x + 5 \sin x - 3 = 0.$$

(2)

(b) Solve, for  $0 \leq x < 360^\circ$ ,

$$2 \sin^2 x + 5 \sin x - 3 = 0.$$

(4)

### Extension

- 1 [MAT 2010 1C] In the range  $0 \leq x < 360^\circ$ , the equation
- $$\sin^2 x + 3 \sin x \cos x + 2 \cos^2 x = 0$$
- Has how many solutions?
- 2 [MAT 2014 1E] As  $x$  varies over the real numbers, the largest value taken by the function
- $$(4 \sin^2 x + 4 \cos x + 1)^2$$
- equals what?